

APPENDIX A:

SUPPLEMENTAL TABLES FOR CHAPTER 2

Table A1. Sampling locations for wild sugar kelp blades, including justification for contaminated site designations. Column ‘#’ represents the corresponding labels in map figures. Counts for ‘n’ indicate number of blades sampled. Two samples were taken from each blade.

Region	#	Site	n	Latitude	Longitude	Nearby contamination
Gloucester Harbor	1	Atlantic Road (ATL)	3	42.6069	-70.6309	Golf course, sewage outfall
	2	Brace Rock (BRK)	5	42.5256	-70.8696	-
	3	Norman’s Woe (NORM)	5	42.5788	-70.6935	-
Salem Harbor	4	Misery Island (MIS)	5	42.5464	-70.7983	-
	5	Baker Island (BK)	5	42.5351	-70.7893	-
	6	Eagle Island (EAG)	5	42.5252	-70.8119	Sewage outfall, coal plant
Marblehead	7	Tinker Island (TKR)	5	42.4811	-70.8334	-
	8	Rams Island (RAM)	5	42.4752	-70.8613	-
	9	Great Pigs Rock (PIG)	2	42.4618	-70.8531	-
Boston Harbor	10	Gallops Island (GAL)	5	42.3276	-70.9442	Heavy urbanization, sewage outfall
Farmed sites	-	Southern Farm (SFARM)	12	<i>anonymous</i>	<i>anonymous</i>	-
	-	Northern Farm (NFARM)	8	<i>anonymous</i>	<i>anonymous</i>	-

Table A2. Analysis of variance table for generalized linear models with Gamma distributions and log links. Heavy metal concentration (ppm) is the response variable to sample site and blade location. Sample size = 117.

Metal	Factor	df	Deviance	Residual Deviance	F Value	Pr(>F)
<i>Total arsenic (AsT)</i>	Site	10	2.05	10.39	5.58	<0.0001
	Blade Location	1	6.53	3.86	178.25	<0.0001
	Site*Blade Location	10	0.65	3.21	1.78	0.08
	Residuals	104	-	12.44	-	-
<i>Inorganic arsenic (iAs)</i>	Site	10	42.76	23.87	43.16	<0.0001
	Blade Location	1	3.80	20.07	38.32	<0.0001
	Site*Blade Location	10	11.52	8.55	11.63	<0.0001
	Residuals	104	-	66.62	-	-
<i>Cadmium (Cd)</i>	Site	10	4.48	73.97	1.81	0.07
	Blade Location	1	50.30	23.67	203.19	<0.0001
	Site*Blade Location	10	3.71	19.96	1.50	0.15
	Residuals	104	-	78.44	-	-
<i>Mercury (Hg)</i>	Site	10	33.36	27.07	13.99	<0.0001
	Blade Location	1	2.29	24.78	9.61	0.003
	Site*Blade Location	10	3.806	20.98	1.60	0.12
	Residuals	104	-	60.43	-	-
<i>Lead (Pb)</i>	Site	10	93.18	52.34	20.62	<0.0001
	Blade Location	1	2.11	50.24	4.66	0.034
	Site*Blade Location	10	15.58	34.66	3.45	0.0008
	Residuals	104	-	145.52	-	-

Table A3a-b. Averages \pm standard deviation for concentrations (ppm) of cadmium (Cd), total arsenic (AsT), and inorganic arsenic (iAs) (panel 3a); as well as lead (Pb) and mercury (Hg) (panel 3b) at each sample site and blade location.

A3a.						
Site	AsT		iAs		Cd	
	Distal Tip	Base	Distal Tip	Base	Distal Tip	Base
Atlantic Road	40.94 \pm 4.65	71.32 \pm 4.23	0.10 \pm 0.03	0.16 \pm 0.04	0.35 \pm 0.22	1.91 \pm 0.11
Brace Rock	37.52 \pm 8.62	50.83 \pm 16.92	0.12 \pm 0.05	0.10 \pm 0.04	0.35 \pm 0.10	0.86 \pm 0.86
Norman's Woe	47.12 \pm 5.14	74.14 \pm 14.86	0.09 \pm 0.03	0.11 \pm 0.04	0.49 \pm 0.22	1.42 \pm 0.72
Misery Island	47.04 \pm 6.46	59.00 \pm 14.60	0.13 \pm 0.02	0.10 \pm 0.04	0.46 \pm 0.17	1.38 \pm 1.16
Bakers Island	44.84 \pm 4.27	76.03 \pm 4.04	0.08 \pm 0.02	0.13 \pm 0.06	0.29 \pm 0.09	1.74 \pm 0.62
Eagle Island	44.09 \pm 7.47	101.12 \pm 17.93	0.06 \pm 0.01	0.08 \pm 0.02	0.29 \pm 0.17	2.40 \pm 0.49
Tinker's Island	62.41 \pm 4.37	36.69 \pm 5.78	0.10 \pm 0.03	0.05 \pm 0.01	0.40 \pm 0.20	1.45 \pm 0.76
Rams Island	47.53 \pm 7.81	80.18 \pm 15.77	0.19 \pm 0.06	0.08 \pm 0.04	0.46 \pm 0.14	1.96 \pm 0.69
Great Pig Rocks	63.41 \pm 14.18	43.94 \pm 9.53	0.14 \pm 0.03	0.09 \pm 0.04	0.46 \pm 0.27	1.41 \pm 0.28
Gallops Island	43.20 \pm 13.78	83.96 \pm 5.69	0.99 \pm 0.41	0.15 \pm 0.03	0.23 \pm 0.17	1.55 \pm 0.44
Northern Farmed	38.45 \pm 6.61	63.50 \pm 15.83	0.26 \pm 0.07	0.11 \pm 0.03	0.27 \pm 0.16	1.39 \pm 0.34
A3b.						
Site	Pb		Hg			
	Distal tip	Base	Distal tip	Base		
Atlantic Road	0.45 \pm 0.31	1.53 \pm 1.47	0.05 \pm 0.01	0.09 \pm 0.06		
Brace Rock	0.29 \pm 0.30	0.28 \pm 0.30	0.05 \pm 0.04	0.03 \pm 0.02		
Norman's Woe	0.17 \pm 0.03	0.24 \pm 0.28	0.09 \pm 0.08	0.05 \pm 0.03		
Misery Island	2.86 \pm 4.05	1.42 \pm 0.63	0.03 \pm 0.01	0.02 \pm 0.01		
Bakers Island	1.75 \pm 0.65	3.89 \pm 2.19	0.03 \pm 0.00	0.02 \pm 0.01		
Eagle Island	0.48 \pm 0.19	0.52 \pm 0.24	0.01 \pm 0.01	0.02 \pm 0.01		
Tinker's Island	0.28 \pm 0.04	0.19 \pm 0.07	0.02 \pm 0.01	0.02 \pm 0.001		
Rams Island	0.93 \pm 0.64	0.46 \pm 0.32	0.04 \pm 0.03	0.02 \pm 0.01		
Great Pig Rocks	0.34 \pm 0.12	0.34 \pm 0.22	0.03 \pm 0.01	0.02 \pm 0.00		
Gallops Island	6.28 \pm 2.76	0.83 \pm 0.16	0.09 \pm 0.02	0.09 \pm 0.03		
Northern Farmed	1.76 \pm 0.64	0.82 \pm 0.38	0.06 \pm 0.02	0.04 \pm 0.01		

Table A4. Post-hoc contrasts of concentrations of heavy metals (ppm) at each site in samples taken from the base blades compared to samples taken at the distal tip of blades. Results are given on the log scale.

Metal	Sample site	estimate	SE	Z-ratio	p-value	
<i>Total arsenic</i>	Atlantic Road**	0.56	0.16	3.55	0.0004	
	Baker's***	0.53	0.13	4.11	<0.0001	
	Brace Rock*	0.30	0.12	2.51	0.01	
	Eagle***	0.83	0.12	6.85	<0.0001	
	Gallops***	0.66	0.12	5.49	<0.0001	
	Misery	0.23	0.12	1.87	0.06	
	North Farmed***	0.50	0.10	5.24	<0.0001	
	Norman's Woe**	0.45	0.12	3.74	0.0002	
	Pigs	0.37	0.19	1.92	0.06	
	Rams***	0.52	0.12	4.32	<0.0001	
	Tinkers***	0.53	0.12	4.39	<0.0001	
	<i>Inorganic arsenic</i>	Atlantic Road*	0.54	0.26	2.09	0.04
		Baker's*	0.45	0.21	2.13	0.03
Brace Rock		-0.21	0.20	-1.07	0.29	
Eagle		0.29	0.20	1.46	0.14	
Gallops***		-1.89	0.20	-9.50	<0.0001	
Misery		-0.26	0.20	-1.31	0.19	
North Farmed***		-0.85	0.16	-5.42	<0.0001	
Norman's Woe		0.16	0.20	0.78	0.434	
Pigs		-0.46	0.32	-1.46	0.15	
Rams***		-0.83	0.20	-4.15	<0.0001	
Tinkers**		-0.74	0.20	-3.74	0.0002	
<i>Cadmium</i>		Atlantic Road***	1.69	0.41	4.15	<0.0001
		Baker's***	1.79	0.33	5.36	<0.0001
	Brace Rock**	0.90	0.32	2.87	0.004	
	Eagle***	2.10	0.32	6.78	<0.0001	
	Gallops***	1.93	0.32	6.12	<0.0001	
	Misery**	1.07	0.32	3.40	0.0007	
	North Farmed***	1.62	0.25	6.53	<0.0001	
	Norman's Woe**	1.07	0.32	3.40	0.0007	
	Pigs*	1.11	0.50	2.23	0.02	
	Rams***	1.41	0.32	4.48	<0.0001	
	Tinkers***	1.23	0.32	4.07	<0.0001	
	<i>Lead</i>	Atlantic Road*	1.21	0.55	2.21	0.03
		Baker's	0.80	0.45	1.77	0.08
Brace Rock		-0.07	0.43	-0.15	0.88	
Eagle		0.08	0.43	0.20	0.85	
Gallops***		-2.02	0.43	-4.75	<0.0001	
Misery		-0.70	0.43	-1.65	0.10	
North Farmed*		-0.76	0.34	-2.27	0.02	
Norman's Woe		0.36	0.43	0.85	0.39	
Pigs		-0.009	0.62	-0.013	0.99	
Rams		-0.70	0.43	-1.64	0.10	
Tinkers		-0.40	0.43	-0.94	0.35	
<i>Mercury</i>		Atlantic Road	0.66	0.40	1.66	0.10
		Baker's	-0.23	0.33	-0.70	0.48
	Brace Rock*	-0.64	0.31	-2.06	0.04	
	Eagle	0.29	0.31	0.95	0.34	
	Gallops	0.03	0.31	0.10	0.92	
	Misery	-0.31	0.31	-1.00	0.32	
	North Farmed	-0.38	0.24	-1.56	0.12	
	Norman's Woe	-0.58	0.31	-1.87	0.06	
	Pigs	-0.49	0.49	-1.00	0.32	
	Rams**	-0.84	0.31	-2.72	0.007	
	Tinkers	-0.47	0.31	-1.51	0.13	

Table A5. Post-hoc contrasts of concentrations of total arsenic (AsT, ppm) in samples taken from different sample sites, but not accounting for blade location. Results are given on the log scale.

Contrast	estimate	SE	Z-ratio	p-value
ATL – BK	-0.08	0.10	-0.77	1.00
ATL – BRK	0.21	0.10	2.15	0.54
ATL – EAG	-0.21	0.10	-2.14	0.55
ATL – GAL	-0.11	0.10	-1.10	0.99
ATL – MIS	0.03	0.10	0.26	1.00
ATL – NFARM	0.09	0.09	0.98	1.00
ATL – NORM	-0.09	0.10	-0.91	1.00
ATL – PIG	0.02	0.12	0.19	1.00
ATL – RAM	-0.13	0.10	-1.35	0.96
ATL – TKR	0.12	0.10	1.23	0.98
BK – BRK *	0.29	0.09	3.29	0.04
BK – EAG	-0.13	0.09	-1.52	0.91
BK – GAL	-0.03	0.09	-0.35	1.00
BK – MIS	0.10	0.09	1.16	0.99
BK – NFARM	0.17	0.08	2.08	0.59
BK – NORM	-0.01	0.09	-0.14	1.00
BK – PIG	0.10	0.12	0.87	1.00
BK – RAM	-0.06	0.09	-0.63	1.00
BK – TKR	0.20	0.09	2.25	0.47
BRK – EAG *	-0.42	0.09	-4.96	<0.0001
BRK – GAL*	-0.32	0.09	-3.75	0.01
BRK – MIS	-0.19	0.09	-2.19	0.51
BRK – NFARM	-0.12	0.08	-1.60	0.88
BRK – NORM*	-0.30	0.09	-3.54	0.02
BRK – PIG	-0.19	0.11	-1.67	0.85
BRK – RAM**	-0.35	0.09	-4.04	0.003
BRK – TKR	-0.09	0.09	-1.07	0.99
EAG – GAL	0.10	0.09	1.21	0.98
EAG – MIS	0.24	0.09	2.77	0.17
EAG – NFARM**	0.30	0.08	3.90	0.005
EAG – NORM	0.12	0.09	1.42	0.94
EAG – PIG	0.24	0.11	2.08	0.60
EAG – RAM	0.08	0.09	0.92	1.00
EAG – TKR**	0.33	0.09	3.89	0.005
GAL – MIS	0.13	0.09	1.56	0.90
GAL – NFARM	0.20	0.08	2.56	0.27
GAL – NORM	0.02	0.09	0.22	1.00
GAL – PIG	0.13	0.11	1.16	0.99
GAL – RAM	-0.02	0.09	-0.29	1.00
GAL – TKR	0.23	0.09	2.69	0.21
MIS – NFARM	0.06	0.08	0.83	1.00
MIS – NORM	-0.12	0.09	-1.34	0.96
MIS – PIG	-0.00	0.11	-0.02	1.00
MIS – RAM	-0.16	0.09	-1.85	0.75
MIS – TKR	0.10	0.09	1.12	0.99
NFARM – NORM	-0.18	0.08	-2.32	0.42
NFARM – PIG	-0.07	0.11	-0.62	1.00
NFARM – RAM	-0.22	0.08	-2.89	0.13
NFARM – TKR	0.03	0.08	0.42	1.00
NORM – PIG	0.11	0.11	1.00	1.00
NORM – RAM	-0.04	0.09	-0.51	1.00
NORM – TKR	0.21	0.09	2.47	0.32
PIG – RAM	-0.16	0.11	-1.38	0.95
PIG – TKR	0.10	0.11	0.87	1.00
RAM – TKR	0.25	0.09	2.98	0.10

Table A6. Average \pm standard deviation (SD) of concentrations (ppm) of heavy metals in our preliminary Southern Farmed samples from 2018, and comparison of those averages to relevant standards. Preliminary samples were only collected from the mid-part of each blade.

	AsT (n=12)	iAs (n=4)	Cd (n=12)	Pb (n=12)	Hg (n=12)
Average \pm SD (ppm)	32 \pm 2	0.0006 \pm 0.0003	0.72 \pm 0.15	0.82 \pm 0.54	0.05 \pm 0.03
% that exceeded MCL	0%	0%	17%	0%	0%
% of MADL	-	-	17.5%	164%	16.6%
% of NSRL	-	0.006%	-	5.5%	-
Grams of dried kelp safe to eat	-	16,666g	5.7g	18.18g or 0.6g	6g

Table A7. Average percent (\pm standard deviation) of inorganic arsenic (iAs) concentration in relation to total arsenic (AsT) concentration for each sample site and blade location.

Site	Base of blade	Distal tip
Gallops Island (GAL)	0.18 \pm 0.03	2.26 \pm 0.69
Great Pig Rock (PIG)	0.14 \pm 0.09	0.34 \pm 0.17
Rams Island (RAM)	0.11 \pm 0.05	0.40 \pm 0.14
Tinkers Island (TKR)	0.08 \pm 0.02	0.28 \pm 0.09
Eagle Island (EAG)	0.08 \pm 0.02	0.14 \pm 0.03
Baker Island (BK)	0.17 \pm 0.08	0.18 \pm 0.03
Misery Island (MIS)	0.19 \pm 0.12	0.28 \pm 0.07
Norman's Woe (NORM)	0.14 \pm 0.03	0.20 \pm 0.06
Brace Rock (BRK)	0.23 \pm 0.16	0.32 \pm 0.15
Atlantic Road (ATL)	0.23 \pm 0.04	0.23 \pm 0.04
Northern Farm (NFARM)	0.20 \pm 0.09	0.70 \pm 0.21

Table A8. Post-hoc contrasts for inorganic arsenic (iAs) concentrations (ppm) for samples taken at different sample sites, but not accounting for blade location. Results are presented on the log scale.

Contrast	estimate	SE	Z-ratio	p-value
ATL – BK	0.21	0.17	1.27	0.97
ATL – BRK	0.15	0.16	0.95	1.00
ATL – EAG*	0.57	0.16	3.48	0.02
ATL – GAL***	-1.12	0.16	-6.91	<0.0001
ATL – MIS	0.11	0.16	0.66	1.00
ATL – NFARM	-0.32	0.15	-2.13	0.56
ATL – NORM	0.24	0.16	1.45	0.94
ATL – PIG	0.13	0.20	0.65	1.00
ATL – RAM	0.02	0.16	0.12	1.00
ATL – TKR**	0.61	0.16	3.73	0.009
BK – BRK	-0.06	0.15	-0.39	1.00
BK – EAG	0.35	0.15	2.44	0.34
BK – GAL***	-1.33	0.15	-9.19	<0.0001
BK – MIS	-0.10	0.15	-0.72	1.00
BK – NFARM**	-0.53	0.13	-4.04	0.003
BK – NORM	0.02	0.15	0.17	1.00
BK – PIG	-0.08	0.19	-0.42	1.00
BK – RAM	-0.19	0.15	-1.31	0.97
BK – TKR	0.40	0.15	2.73	0.19
BRK – EAG	0.41	0.14	2.92	0.12
BRK – GAL***	-1.28	0.14	-9.07	<0.0001
BRK – MIS	-0.05	0.14	-0.34	1.00
BRK – NFARM**	-0.47	0.13	-3.74	0.009
BRK – NORM	0.08	0.14	0.58	1.00
BRK – PIG	-0.02	0.19	-0.12	1.00
BRK – RAM	-0.13	0.14	-0.95	1.00
BRK – TKR	0.45	0.14	3.22	0.05
EAG – GAL***	-1.69	0.14	-11.99	<0.0001
EAG – MIS*	-0.46	0.14	-3.26	0.044
EAG – NFARM***	-0.89	0.13	-6.98	<0.0001
EAG – NORM	-0.33	0.14	-2.34	0.40
EAG – PIG	-0.43	0.19	-2.33	0.41
EAG – RAM**	-0.54	0.14	-3.87	0.005
EAG – TKR	0.04	0.14	0.30	1.00
GAL – MIS***	1.23	0.14	8.73	<0.0001
GAL – NFARM***	0.80	0.13	6.32	<0.0001
GAL – NORM***	1.36	0.14	9.65	<0.0001
GAL – PIG***	1.25	0.19	6.73	<0.0001
GAL – RAM***	1.14	0.14	8.12	<0.0001
GAL – TKR***	1.73	0.14	12.29	<0.0001
MIS – NFARM*	-0.43	0.13	-3.37	0.031
MIS – NORM	0.13	0.14	0.92	1.00
MIS – PIG	0.03	0.19	0.13	1.00
MIS – RAM	-0.09	0.14	-0.61	1.00
MIS – TKR*	0.50	0.14	3.56	0.017
NFARM – NORM**	0.56	0.13	4.38	0.0006
NFARM – PIG	0.45	0.18	2.57	0.26
NFARM – RAM	0.34	0.13	2.69	0.20
NFARM – TKR***	0.93	0.13	7.31	<0.0001
NORM – PIG	-0.10	0.19	-0.56	1.00
NORM – RAM	-0.22	0.14	-1.53	0.91
NORM – TKR	0.37	0.14	2.64	0.23
PIG – RAM	-0.11	0.19	-0.60	1.00
PIG – TKR	0.48	0.19	2.55	0.27
RAM – TKR**	0.59	0.14	4.17	0.002

Table A9. Post-hoc contrasts for lead (Pb) concentrations (ppm) for samples taken at different sample sites, but not accounting for blade location. Results are presented on the log scale.

Contrast	estimate	SE	Z-ratio	p-value
ATL – BK	-1.15	0.36	-3.22	0.05
ATL – BRK	1.07	0.35	3.08	0.07
ATL – EAG	0.50	0.35	1.45	0.93
ATL – GAL	-1.01	0.35	-2.91	0.12
ATL – MIS	-0.89	0.35	-2.55	0.27
ATL – NFARM	-0.37	0.32	-1.14	0.99
ATL – NORM**	1.42	0.35	4.10	0.002
ATL – PIG	0.90	0.43	2.07	0.60
ATL – RAM	0.24	0.35	0.68	1.00
ATL – TKR*	1.28	0.35	3.68	0.01
BK – BRK***	2.21	0.31	7.15	<0.0001
BK – EAG***	1.65	0.31	5.32	<0.0001
BK – GAL	0.13	0.31	0.43	1.00
BK – MIS	0.26	0.31	0.84	1.00
BK – NFARM	0.78	0.28	2.77	0.17
BK – NORM***	2.57	0.31	8.29	<0.0001
BK – PIG***	2.04	0.40	5.05	<0.0001
BK – RAM**	1.38	0.31	4.46	0.0004
BK – TKR***	2.42	0.31	7.82	<0.0001
BRK – EAG	-0.57	0.30	-1.88	0.73
BRK – GAL***	-2.08	0.30	-6.92	<0.0001
BRK – MIS***	-1.96	0.30	-6.50	<0.0001
BRK – NFARM***	-1.44	0.27	-5.30	<0.0001
BRK – NORM	0.35	0.30	1.18	0.99
BRK – PIG	-0.17	0.40	-0.43	1.00
BRK – RAM	-0.83	0.30	-2.78	0.17
BRK – TKR	0.21	0.30	0.69	1.00
EAG – GAL***	-1.52	0.30	-5.04	<0.0001
EAG – MIS**	-1.39	0.30	-4.62	0.002
EAG – NFARM	-0.87	0.27	-3.21	0.05
EAG – NORM	0.92	0.30	3.06	0.08
EAG – PIG	0.39	0.40	0.99	1.00
EAG – RAM	-0.27	0.30	-0.89	1.00
EAG – TKR	0.77	0.30	2.57	0.27
GAL – MIS	0.13	0.30	0.42	1.00
GAL – NFARM	0.64	0.27	2.38	0.38
GAL – NORM***	2.43	0.30	8.10	<0.0001
GAL – PIG***	1.91	0.40	4.80	0.0001
GAL – RAM**	1.25	0.30	4.15	0.002
GAL – TKR***	2.29	0.30	7.61	<0.0001
MIS – NFARM	0.52	0.27	1.92	0.71
MIS – NORM***	2.31	0.30	7.68	<0.0001
MIS – PIG**	1.78	0.40	4.49	0.0004
MIS – RAM**	1.12	0.30	3.73	0.009
MIS – TKR***	2.16	0.30	7.19	<0.0001
NFARM – NORM***	1.79	0.27	6.60	<0.0001
NFARM – PIG*	1.27	0.38	3.37	0.03
NFARM – RAM	0.60	0.27	2.22	0.49
NFARM – TKR***	1.64	0.27	6.06	<0.0001
NORM – PIG	-0.52	0.40	-1.32	0.97
NORM – RAM**	-1.19	0.30	-3.95	0.004
NORM – TKR	-0.15	0.30	-0.49	1.00
PIG – RAM	-0.66	0.40	-1.67	0.85
PIG – TKR	0.38	0.40	0.95	1.00
RAM – TKR*	1.04	0.30	3.46	0.023

Table A10. Post-hoc contrasts of lead (Pb) concentration (ppm) in samples taken from the base of sugar kelp blades at different sample sites. Results are given on the log scale.

BASE OF BLADE				
Contrast	estimate	SE	Z-ratio	p-value
ATL – BK	-0.94	0.49	-1.91	0.71
ATL – BRK*	1.71	0.49	3.48	0.02
ATL – EAG	1.07	0.49	2.18	0.52
ATL – GAL	0.61	0.49	1.23	0.98
ATL – MIS	0.07	0.49	0.15	1.00
ATL – NFARM	0.62	0.46	1.37	0.96
ATL – NORM*	1.85	0.49	3.77	0.01
ATL – PIG	1.51	0.61	2.46	0.33
ATL – RAM	1.19	0.49	2.42	0.35
ATL – TKR**	2.08	0.49	4.24	0.001
BK – BRK***	2.65	0.43	6.22	<0.0001
BK – EAG***	2.01	0.43	4.72	0.0001
BK – GAL*	1.54	0.43	3.63	0.01
BK – MIS	1.01	0.43	2.37	0.38
BK – NFARM**	1.56	0.38	4.07	0.002
BK – NORM***	2.78	0.43	6.55	<0.0001
BK – PIG**	2.45	0.56	4.35	0.0007
BK – RAM***	2.13	0.43	5.00	<0.0001
BK – TKR***	3.02	0.43	7.10	<0.0001
BRK – EAG	-0.64	0.43	-1.50	0.92
BRK – GAL	-1.10	0.43	-2.59	0.25
BRK – MIS**	-1.64	0.43	-3.85	0.005
BRK – NFARM	-1.09	0.38	-2.84	0.14
BRK – NORM	0.14	0.43	0.33	1.00
BRK – PIG	-0.20	0.56	-0.35	1.00
BRK – RAM	-0.52	0.43	-1.22	0.98
BRK – TKR	0.37	0.43	0.88	1.00
EAG – GAL	-0.46	0.43	-1.09	0.99
EAG – MIS	-1.00	0.43	-2.35	0.40
EAG – NFARM	-0.45	0.38	-1.17	0.99
EAG – NORM	0.78	0.43	1.83	0.76
EAG – PIG	0.44	0.56	0.78	1.00
EAG – RAM	0.12	0.43	0.28	1.00
EAG – TKR	1.01	0.43	2.38	0.38
GAL – MIS	-0.53	0.43	-1.26	0.98
GAL – NFARM	0.02	0.38	0.04	1.00
GAL – NORM	1.24	0.43	2.92	0.12
GAL – PIG	0.90	0.56	1.61	0.88
GAL – RAM	0.58	0.43	1.37	0.95
GAL – TKR*	1.48	0.43	3.47	0.02
MIS – NFARM	0.55	0.38	1.44	0.94
MIS – NORM**	1.78	0.43	4.18	0.002
MIS – PIG	1.44	0.56	2.56	0.27
MIS – RAM	1.12	0.43	2.63	0.23
MIS – TKR***	2.01	0.43	4.73	0.0001
NFARM – NORM	1.23	0.38	3.20	0.05
NFARM – PIG	0.89	0.53	1.67	0.85
NFARM – RAM	0.57	0.38	1.48	0.93
NFARM – TKR**	1.46	0.38	3.81	0.007
NORM – PIG	-0.34	0.56	-0.60	1.00
NORM – RAM	-0.66	0.43	-1.55	0.90
NORM – TKR	0.24	0.43	0.55	1.00
PIG – RAM	-0.32	0.56	-0.57	1.00
PIG – TKR	0.57	0.56	1.02	1.00
RAM – TKR	0.89	0.43	2.10	0.58

Table A11. Post-hoc contrasts of lead (Pb) concentration (ppm) in samples taken from the distal tip of sugar kelps blades at different sample sites. Results are given on the log scale.

DISTAL TIP OF BLADE				
Contrast	estimate	SE	Z-ratio	p-value
ATL – BK	-1.35	0.51	-2.64	0.23
ATL – BRK	0.43	0.49	0.88	1.00
ATL – EAG	-0.06	0.49	-0.12	1.00
ATL – GAL	-2.63	0.49	-5.35	0.00
ATL – MIS	-1.84	0.49	-3.75	0.01
ATL – NFARM	-1.35	0.46	-2.98	0.10
ATL – NORM	1.00	0.49	2.03	0.63
ATL – PIG	0.29	0.61	0.47	1.00
ATL – RAM	-0.72	0.49	-1.47	0.93
ATL – TKR	0.47	0.49	0.96	1.00
BK – BRK *	1.78	0.45	3.96	0.00
BK – EAG	1.29	0.45	2.87	0.13
BK – GAL	-1.27	0.45	-2.83	0.15
BK – MIS	-0.49	0.45	-1.09	0.99
BK – NFARM	-0.00	0.41	-0.00	1.00
BK – NORM	2.35	0.45	5.21	0.00
BK – PIG	1.64	0.58	2.82	0.15
BK – RAM	0.63	0.45	1.41	0.95
BK – TKR	1.82	0.45	4.04	0.00
BRK – EAG *	-0.49	0.43	-1.16	0.99
BRK – GAL*	-3.06	0.43	-7.20	0.00
BRK – MIS	-2.27	0.43	-5.35	0.00
BRK – NFARM	-1.79	0.38	-4.66	0.00
BRK – NORM*	0.57	0.43	1.33	0.96
BRK – PIG	-0.14	0.56	-0.25	1.00
BRK – RAM**	-1.15	0.43	-2.70	0.20
BRK – TKR	0.04	0.43	0.09	1.00
EAG – GAL	-2.57	0.43	-6.04	0.00
EAG – MIS	-1.78	0.43	-4.19	0.00
EAG – NFARM**	-1.29	0.38	-3.38	0.03
EAG – NORM	1.06	0.43	2.49	0.31
EAG – PIG	0.35	0.56	0.62	1.00
EAG – RAM	-0.66	0.43	-1.55	0.90
EAG – TKR**	0.53	0.43	1.25	0.98
GAL – MIS	0.79	0.43	1.85	0.75
GAL – NFARM	1.27	0.38	3.32	0.04
GAL – NORM	3.63	0.43	8.53	0.00
GAL – PIG	2.92	0.56	5.19	0.00
GAL – RAM	1.91	0.43	4.49	0.00
GAL – TKR	3.10	0.43	7.29	0.00
MIS – NFARM	0.49	0.38	1.27	0.97
MIS – NORM	2.84	0.43	6.68	0.00
MIS – PIG	2.13	0.56	3.79	0.01
MIS – RAM	1.12	0.43	2.64	0.23
MIS – TKR	2.31	0.43	5.44	0.00
NFARM – NORM	2.35	0.38	6.14	0.00
NFARM – PIG	1.64	0.53	3.09	0.07
NFARM – RAM	0.64	0.38	1.66	0.86
NFARM – TKR	1.82	0.38	4.76	0.00
NORM – PIG	-0.71	0.56	-1.26	0.97
NORM – RAM	-1.72	0.43	-4.04	0.00
NORM – TKR	-0.53	0.43	-1.24	0.98
PIG – RAM	-1.01	0.56	-1.79	0.79
PIG – TKR	0.18	0.56	0.32	1.00
RAM – TKR	1.19	0.43	2.80	0.16

Table A12. Post-hoc contrasts for mercury (Hg) concentrations (ppm) for samples taken at different sample sites, but not accounting for blade location. Results are presented on the log scale.

Contrast	estimate	SE	Z-ratio	p-value
ATL – BK**	1.13	0.26	4.40	0.0006
ATL – BRK	0.60	0.25	2.36	0.39
ATL – EAG***	1.53	0.25	6.08	<0.0001
ATL – GAL	-0.31	0.25	-1.22	0.98
ATL – MIS	0.96	0.25	3.82	0.01
ATL – NFARM*	0.30	0.23	1.27	0.97
ATL – NORM	-0.02	0.25	-0.07	1.00
ATL – PIG*	1.04	0.32	3.30	0.04
ATL – RAM	0.82	0.25	3.25	0.05
ATL – TKR***	1.25	0.25	4.96	<0.0001
BK – BRK	-0.54	0.23	-2.39	0.37
BK – EAG	0.40	0.23	1.77	0.80
BK – GAL***	-1.44	0.23	-6.40	<0.0001
BK – MIS	-0.17	0.23	-0.76	1.00
BK – NFARM**	-0.84	0.20	-4.10	0.002
BK – NORM*	-1.15	0.23	-5.12	<0.0001
BK – PIG	-0.10	0.29	-0.32	1.00
BK – RAM	-0.32	0.23	-1.40	0.95
BK – TKR	0.12	0.23	0.52	1.00
BRK – EAG**	0.94	0.22	4.29	0.001
BRK – GAL**	-0.90	0.22	-4.14	0.002
BRK – MIS	0.37	0.22	1.68	0.85
BRK – NFARM	-0.30	0.20	-1.52	0.91
BRK – NORM	-0.62	0.22	-2.82	0.15
BRK – PIG	0.44	0.29	1.53	0.91
BRK – RAM	0.22	0.22	1.02	1.00
BRK – TKR	0.65	0.22	3.00	0.09
EAG – GAL***	-1.84	0.22	-8.42	<0.0001
EAG – MIS	-0.57	0.22	-2.61	0.24
EAG – NFARM***	-1.24	0.20	-6.28	<0.0001
EAG – NORM***	-1.55	0.22	-7.10	<0.0001
EAG – PIG	-0.49	0.29	-1.71	0.83
EAG – RAM*	-0.71	0.22	-3.27	0.04
EAG – TKR	-0.28	0.22	-1.29	0.97
GAL – MIS***	1.27	0.22	5.82	<0.0001
GAL – NFARM	0.60	0.20	3.07	0.08
GAL – NORM	0.29	0.22	1.32	0.97
GAL – PIG**	1.35	0.29	4.66	0.0002
GAL – RAM***	1.13	0.22	5.15	<0.0001
GAL – TKR***	1.56	0.22	7.14	<0.0001
MIS – NFARM*	-0.67	0.20	-3.38	0.03
MIS – NORM**	-0.98	0.22	-4.49	0.0004
MIS – PIG	0.08	0.29	0.26	1.00
MIS – RAM	-0.14	0.22	-0.66	1.00
MIS – TKR	0.29	0.22	1.32	0.97
NFARM – NORM	-0.32	0.20	-1.60	0.88
NFARM – PIG	0.74	0.27	2.72	0.19
NFARM – RAM	0.52	0.20	2.65	0.22
NFARM – TKR***	0.95	0.20	4.85	0.0001
NORM – PIG**	1.06	0.29	3.66	0.006
NORM – RAM**	0.84	0.22	3.83	0.006
NORM – TKR***	1.27	0.22	5.82	<0.0001
PIG – RAM	-0.22	0.29	-0.76	1.00
PIG – TKR	0.21	0.29	0.73	1.00
RAM – TKR	0.43	0.22	1.98	0.66